

**Lake Characteristics:**

**Location:** Boom Lake, Rhinelander, Wisconsin, USA

**Access:** Hodag Park boat landing. Large paved parking lot with improved lake access. Minutes from the lodging.

**Maximum Depth:** 10 meters

**Median Depth:** 3 meters

**Water Source:** Boom Lake is an impoundment of the Wisconsin River. The Rhinelander Paper Company maintains the water level.

**Shoreline:** Nearly entirely upland and fairly well developed. There are a number of islands and numerous bays.


**Bottom:** The bottom is comprised mostly of muck and sand. The bays and backwaters are largely muck. The Southeast side along the park has a diverse bottom of rock, sand, and muck.

**Water:** Moderately fertile and medium brown in color. Good water quality.

**Vegetation:** There is not an abundance of vegetation in the main basin, however, the shallow bays and smaller basins have moderated vegetation. Primary weed types are elodea, coontail, milfoil, pondweed, duckweed, arrowhead, and lily pad.

**Fishery**

**Species:** White Crappie (*Pomoxis annularis*), Black Crappie (*Pomoxis nigromaculatus*), Bluegill (*Lepomis macrochirus*), Pumpkinseed (*Lepomis gibbosus*), Perch (*Perca flavescens*), Yellow Bullhead (*Ictalurus natalis*), White Sucker (*Catostomus commersonii*), Creek Chub (*Semotilus atromaculatus*), Golden Shiner (*Notemigonus crysoleucas*)



**Lake Characteristics:**

**Location:** Lake George, Rhinelander, Wisconsin, USA

**Access:** Boat landing, with dock and parking on the East shore.

**Maximum Depth:** 7.9 meters

**Median Depth:** 3 meters

**Water Source:** Lake George is a drainage lake

**Shoreline:** Nearly entirely upland and fairly well developed.


**Bottom:** The bottom is comprised of a mix of muck, sand, gravel, and rocks.

**Water:** Moderately fertile and light brown in color

**Vegetation:** The water is fertile enough to sustain a good population of submergent vegetation, including cabbage weed, coontail, duckweed, pondweed, bulrush, and lily pads. There are scattered weedbeds along the lakes shorelines.

**Fishery**

**Species:** White Crappie (*Pomoxis annularis*), Black Crappie (*Pomoxis nigromaculatus*), Bluegill (*Lepomis macrochirus*), Pumpkinseed (*Lepomis gibbosus*), Perch (*Perca flavescens*), Yellow Bullhead (*Ictalurus natalis*), White Sucker (*Catostomus commersonii*), Creek Chub (*Semotilus atromaculatus*), Golden Shiner (*Notemigonus crysoleucas*)



### Lake Characteristics:

**Location:** Lake Thompson, Rhinelander, Wisconsin, USA

**Access:** Boat landing with roadside parking off North Shore Drive.

**Maximum Depth:** 10 meters

**Median Depth:** 3 meters

**Water Source:** Lake Thompson is a drainage lake

**Shoreline:** Nearly entirely upland and fairly well developed.

**Bottom:** The bottom is comprised of a mix of muck, sand, gravel, and rubble.

**Water:** Moderately fertile and light brown in color

**Vegetation:** There are long stretches of shoreline that contain bulrushes on Thompson. The outer edge of the rushes, run into a mixture of cabbage and coontail. There are also lily pads, pondweed, and duckweed. The weed edge is shallow, due to the water clarity. The outside weed edge is only 4 to 6 feet on average around the lake.

### Fishery

**Species:** White Crappie (*Pomoxis annularis*), Black Crappie (*Pomoxis nigromaculatus*), Bluegill (*Lepomis macrochirus*), Pumpkinseed (*Lepomis gibbosus*), Perch (*Perca flavescens*), Yellow Bullhead (*Ictalurus natalis*), White Sucker (*Catostomus commersonii*), Creek Chub (*Semotilus atromaculatus*), Golden Shiner (*Notemigonus crysoleucas*)

